COMPTE
RENDU
Vol.2

G E S C O

"Souches" de Van Gogh

Montpellier - 3 au 7 JUILLET 2001

VILLE DE FUISSEQUIER

INSTITUT SUPERIEUR
DE LA VIGNE ET DU VIN

Ecole Nationale Supérieure d'Agriculture

PARTIAL ROOT-DRYING: EFFECTS ON THE PHYSIOLOGY AND PRODUCTION IN GRAPEVINE

DESSÈCHEMENT PARTIEL DES RACINES: EFFETS SUR LA PHYSIOLOGIE ET LA PRODUCTION DE LA VIGNE

C.R. SOUZA (2); T. SANTOS (1), J.P. MAROCO (2), M.L. RODRIGUES (1), C. LOPES (1), J.S. PEREIRA (1), M.M. CHAVES (1,2)

⁽¹⁾ Instituto Superior de Agronomia, Lisboa.

⁽²⁾ ITQB, Oeiras. Portugal.

1. INTRODUCTION

The objective of the present work was to evaluate a system of partial root drying irrigation on grapevine physiology and productivity as compared to traditional systems of irrigation. In Partial Root Drying (PRD) irrigation, water is applied in alternate rows of the vine, allowing for part of the root system to be always in contact with drying soil, while the other half is in contact with wet soil. We hypothesize that with a PRD system we can increase water use efficiency by limiting the vegetative vigour, without reducing the amount and the quality of the berries. Vine responses to soil water availability under the different irrigation schemes were evaluated in terms of gas exchange, water relations (sap flow and leaf water potential) and production.

2. MATERIAL & METHODS

Experiments were conducted in 4-year old *Vitis vinifera* cv. Muscat of Alexandria grafted in rootstock 1.103 Paulsen growing in an experimental vineyard belonging to the "Centro Experimental de Pegões". The climate of the region is characterised by hot and dry summers. The soil is derived from podzols with sand in the surface layers (0.6-1.0 m) and clay-like in the deepest layers. The irrigation treatments were: non irrigated but rain-fed (NI); partial root drying (PRD, 50% of the ETc was supplied to only one side of the root system, alternating sides each 15 days); full irrigated (FI, 100 % of the ETc was supplied to both the sides of the root system); deficit irrigated (DI, 50% of the ETc was supplied to both sides of the row. Measurements of net CO₂ assimilation (A) and stomatal conductance (gs), were done weekly, from the beginning of berry development until harvest, using an IRGA (LI-COR, model LI-6400). Full mature, sun-exposed leaves, were measured at 05:00, 10:00, 12:00, 14:00, 16:00 and 18:00h (local time) with 4 replications per treatment. Chlorophyll a fluorescence was measured using a Mini Pam fluorometer (H.Walz), in the same leaves used for gas-exchange measurements. Sap flow measurements were done with a sap flow measurement system (Flow 32, Dynamax) installed in main branches of the vines. Leaf water potential (ψ_L) was measured with a Scholander-type pressure chamber (PMS). Berry maturation was followed periodically until harvest, by sampling 200 berries per treatment as proposed by Carbonneau (1991).

3. RESULTS & DISCUSSION

Pre-dawn water potential decreased in NI vines from July onward, being significantly lower than ψ_L in PRD, DI, and FI vines (which showed an almost constant ψ_L , always higher than -0.2Mpa, throughout the growing season). By the end of the season ψ_L was significantly lower in DI than in PRD vines, these being lower than ψ_L of FI vines (Fig.1). NI vines showed the largest reductions in stomatal conductance and photosynthesis, reaching values of 0.10 mol.m⁻²s⁻¹ (± 0.02) and 5 μ molm⁻²s⁻¹ (± 1.15), respectively, by the end of August. FI vines exhibited the largest values, with A between 14 to 12 μ molm⁻²s⁻¹ and gs between 0.3 and 0.5 molm⁻²s⁻¹ as shown in the A and g course for one typical day (Fig. 2B). Statistically significant differences in photosynthesis and stomatal conductance were only observed between the watered treatments (FI, PRD and DI) and the non-watered (NI). However, there were significant difference between FI and DI vs. PRD and NI when measurements of A and gs were done under controlled CO₂, light and temperature conditions (Fig 2A). Sap flow measurements showed differences between all watering treatments on clear days, but not on over-cloud days when transpiration demand was lower (Fig.3). These results suggest that PRD vines kept stomata more closed and were able to reduce transpiration losses more substantially than DI vines, in spite of the same amount of water received. No photoinhibition was observed, as demonstrated by pre-dawn Fv/Fm being always above 0.7 in all treatments, during the growing season. The quantum yields decreased only in NI suggesting a down regulation of PSII centers.

Yield was significantly higher in the FI treatment as compared with PRD, DI and NI treatments (table 1), although berry weight was higher in PRD and DI than in NI. Irrigation effects were more pronounced in total phenols ,

which decreased in DI and FI, but not in PRD, than in sugar and acidity. In conclusion, PRD reduced significantly the amount of water used by the plant without reducing berry quality and with only a marginal reduction of yield, as compared with FI.

Figure 1. Pre-dawn leaf water potential in FI, PRD, DI and NI vines. Values are means of 4 replicates, affected by the standard error

Figure 2. Panel A: Photosynthesis (A) and stomatal conductance (g_s) in cv. Muscat of Alexandria on 08/22/00 under constant conditions - 350 ppm CO_2 , 1200 $\mu mol m^{-2} s^{-1}$ PPFD, 25 °C leaf temperature. Panel B: Diurnal gas-exchange course on a typical day. Values are means of 4 replicates, affected by the SE.

Figure 3. Sap flow rates in FI, PRD, DI and NI vines.

Table 1. Yield, berry weight and quality parameters for FI, DI, PRD and NI treatments. Different letter suffixes show statistically significant differences at $p=0.05$.

Treatment	Weight of 100 berries (g)	Yield (kg/grapevine)	Yield (t/ha)	Sugar (g/l)	Acidity (g tart ac./l)	Total phenols (IFC)
NI	413,7 a	7,4 a	29,8 a	203,4 a	3,5 a	15,6 a
PRD	555,8 b	7,7a	30,9 a	212,9 a	3,6 a	15,8 a
DI	594,5 b	7,9 ab	31,6 ab	198,2 a	3,8 ab	13,0 b
FI	564,9 b	9,0 b	36,0 b	198,2 a	3,9 b	12,8 b

REFERENCE

CARBONNEAU, A. (1991). Riv. Vitic. Enol. 4: 345-352.